



Triple Threat: “Unpacking The Intersections Of Gender, Neurodivergence, And Disability In Mental Health”

Sona Panamkunnil *

Higher Secondary School Teacher, English. GHSS Thirumeni, Kannur Kerala, India. Email: sonapk226@gmail.com

Article History

Received Date: 06/02/2026

Revised Date: 12/02/2026

Accepted Date: 20 /02/2026

Published Date: 27/02/2026

Abstract

This paper explores the complex intersections of gender, neurodivergence, and disability in shaping mental health experiences. It highlights the challenges and opportunities in addressing mental health, including stigma, treatment gaps, and accessibility barriers. The interplay of these factors significantly impacts mental health outcomes, with societal gender expectations, diagnostic biases, and accommodation gaps exacerbating disparities. Neurodivergent individuals and those with disabilities face unique mental health challenges, often compounded by socioeconomic inequalities.

The study examines how intersectional identities influence mental health support access and treatment efficacy. It underscores the need for inclusive, culturally sensitive approaches that address the social and medical models of disability. By unpacking these intersections, this research aims to contribute to a deeper understanding of mental health as a basic human right, crucial for personal, community, and socio-economic development. The findings emphasize the importance of tailored interventions, policy reforms, and awareness initiatives to promote mental health equity and social justice.

Keywords:

- Mental health disparities
- Gender
- Neurodivergence
- Disability
- Intersectionality

1. Introduction

The intersection of gender, neurodivergence, and disability significantly shapes mental health outcomes. Mental health is a fundamental human right and a cornerstone of personal well-being, community stability, and socio-economic development. It encompasses our emotional, psychological, and social well-being — influencing how we think, feel, act, and relate to others. In many ways, mental health functions like the software of the brain: when it operates well, life feels manageable; when disrupted, daily functioning becomes significantly more difficult.

However, mental health does not exist in isolation. Social structures, discrimination, economic systems, and cultural expectations deeply influence who receives care, who is believed, and who is left behind.

1.1 Mental Health Challenges and Structural Barriers

Stigma and Discrimination

Stigma remains one of the most significant barriers to mental health care. It creates fear of judgment, social rejection, and discrimination in employment, education, and relationships.

Types of stigma include:

- **Public stigma** – Societal stereotypes (e.g., people with mental illness are violent or burdensome).
- **Self-stigma** – Internalizing negative societal beliefs, leading to shame and low self-esteem.
- **Institutional stigma** – Systemic discrimination such as inadequate insurance coverage or limited services.

Stigma contributes directly to the **treatment gap** by:

- Delaying help-seeking (fear of being labeled or ostracized, stops people from seeking care)
- Reducing treatment adherence (stigma can make people reluctant to stick with treatment)
- Increasing social isolation (reduced quality of life)
- Worsening symptoms (lack of knowledge)

Globally, over 70% of individuals with mental health conditions do not receive adequate treatment, despite many conditions being treatable.

1.2 Shortage of Mental Health Professionals

A global shortage of mental health professionals further widens this gap. In India, for example, there are approximately **0.75 psychiatrists per 100,000 people**, far below the recommended ratio of 3 per 100,000.

Key challenges include:

- Rural-urban disparities (rural areas face a more acute shortage compared to urban areas)
- Burnout among professionals (existing staff are overworked with some having to take on higher patient volumes)
- High demand exceeding workforce capacity. (the need for services outpaced the available workforce, even with government initiatives.)
- Limited postgraduate training seats (there is a need to increase the number of postgraduate seats for psychiatry and clinical psychology.)
- Inadequate training in disability-inclusive care (many healthcare providers, despite training, feel incompetent to handle mental health issues)

Government initiatives such as the National Mental Health Programme (NMHP) and tele-consultation services like Tele-MANAS attempt to address these shortages, but structural gaps remain significant.

1.3 Economic Burden of Mental Health

Mental health conditions carry enormous economic costs. Globally, losses due to mental disorders were estimated at \$1.6 trillion in 2019, projected to rise to \$3–6 trillion by 2030.

Depression and anxiety alone cost approximately \$1 trillion annually in lost productivity. The economic burden includes:

- **Direct costs** – Hospitalization, medication, therapy
- **Indirect costs** – Lost wages, absenteeism, reduced productivity
- **Intangible costs** – Psychological suffering and reduced quality of life

In India, mental health economic losses are projected to exceed \$1 trillion between 2012–2030. Despite this, mental health often receives less than 1% of national health budgets in many regions.

2. Urban–Rural Divide in Mental Health

Urban and rural populations face different yet equally critical mental health challenges. The urban rural mental health divide is characterized by a higher prevalence of mental disorders in urban areas (13.5%) compared to rural areas (6.9%) contrasted by a severe 70-90% treatment gap in rural regions due to lack of professionals, stigmas, and poor infrastructure. Urbanites face higher stress, while rural populations often battle substance abuse and lack of access to care. Urban areas report nearly double the rate of mental disorders (13.5%) compared to rural areas (6.9%) with high rates of depression, anxiety and schizophrenia.

- Higher reported prevalence of depression and anxiety
- Environmental stressors (crowding, pollution, work pressure)
- Social isolation despite population density

Rural Challenges

- Severe shortage of professionals
- Deeply entrenched stigma
- Higher untreated psychosis duration
- Poverty and agricultural stressors
- Limited infrastructure and digital access

While urban areas report higher diagnoses, rural areas experience a “hidden crisis” due to limited detection and treatment.

2.1 Gender and Mental Health

Gender expectations significantly influence mental health outcomes. Women frequently experience higher rates of anxiety and depression due to pressures regarding caregiving, domestic labor and societal norms. Men often face pressure to hide emotions leading to lower help seeking behaviors and higher rates of suicide.

Women experience higher rates of anxiety and depression due to:

- Caregiving expectations
- Domestic labor burden
- Workplace discrimination
- Gender-based violence

They are also 54% more likely to receive psychotropic medication. However, their concerns are often dismissed or labeled as “emotional.”

Men

Men are socialized to suppress vulnerability. This leads to:

- Lower help-seeking behavior
- Higher substance use
- Suicide rates four times higher than women

Although men report fewer diagnoses, they have significantly higher rates of completed suicide — indicating under-detection and internalized distress.

2.2 Gender Non-Conforming and LGBTQ+ Individuals

Individuals who do not confirm to traditional gender roles often face social exclusion, discrimination and stigma which can severely impact their psychological well-being.

2.3 Intersection of Gender & Society

Cultural factors, rather than just biological ones play a major role in these difference, with socialization process heavily influencing how men and women express, or suppress their emotions. Addressing these such as through flexible parental leave and dismantling strict stereotypes is crucial for promoting mental wellness across all genders.

2.4 Gender Disparities in mental health diagnosis & treatment

Women experience higher rates of depression and anxiety often accessing more while men suffer from higher rates of substance abuse and due to lower help seeking have suicide death rates four times higher Gender bias impacts diagnostics with women over diagnosed with depression and men under diagnosed.

Prevalence and Type of Disorder: Women generally have a higher prevalence of internalizing disorders (depression, anxiety, PTSD) while men show higher rates of externalizing disorders (substance abuse, personality disorders)

Diagnostic Bias: Women are more likely to be diagnosed with depression even when reporting identical symptoms to men conversely; men's mental health issues are often under recognized.

Help seeking Behavior: Women are more likely to seek help & receive mental health services often through primary care, compared to men who are less likely to engage with the health system and often do so after severe crises.

Treatment Disparities: Women are 54% more likely to receive psychotropic medication for anxiety and depression.

Suicide rates: Men have significantly higher rates of completed suicide (four times higher than women) despite women having higher rates of attempted suicide and ideation.

Root causes: Disparities are driven by a combination of societal factors (gender stereotypes, stigma), Socioeconomic status, and for women, high rates of gender based or sexual violence.

Healthcare experience – Women frequently report not feeling listened to or taken seriously in clinical interactions (Cambridge Universities press & Assessment.)

Biological Factors: Differences in how men and women metabolize medication can affect treatment efficacy and side effect.

Barriers for men: Societal norms and stigma surrounding mental health often discourage men from seeking help. (National Institutes of Health (NIH))

2.5 LGBTQ+ gender identities & mental health nuances

LGBTQ+ individuals experience elevated rates of depression, anxiety, and suicidality — not due to identity itself, but because of discrimination, minority stress, and social rejection. These mental health challenges are often intensified by marginalization including family rejection bullying and face of discrimination in healthcare particularly for transgender and gender-diverse (TGD) Youth.

Minority Stress Model: LGBTQ+ individuals encounter unique stressors related to their marginalized status, such as internalized stigma, prejudice, and fear of rejection.

Intersectionality: The intersection of sexual orientation or gender identity with other marginalized identities (eg. socioeconomic status, disability) creates a “compounded burden” of stress.

Coming out and concealment: disclosure of one's identity can be a liberating positive experience for mental health, but for many, it carries risks of rejection, causing high anxiety. Conversely constant concealment of identity to avoid, discrimination, can lead to chronic stress, isolation and in some cases severe mental health outcomes.

Healthcare disparities: Many LGBTQ+ Individuals avoid seeking necessary care due to past negative experiences, fear of judgment, or lack of competent, affirmative providers.

2.6 Transgender and Gender-Diverse (TGD) Youth

TGD Youth faces specific, profound challenges including,

- Higher rates of bullying
- Gender dysphoria
- Increased suicide attempts
- Healthcare discrimination

Affirmative care, family support, and community inclusion significantly buffer these risks.

2.7 Neurodivergence and Mental Health

Neurodivergence is a term used to acknowledge that everyone has different brains. This can manifest in many ways, perhaps in how people might process information, communicate behave or function day to day. Neurodivergence recognizes that neurological differences are natural variations in human cognition. Many neurodivergent creative can face barriers in the arts due to complex application forms, rigid communication styles and disruptive working environments. By embedding more accessible and flexible ways of working, you can begin to build a more inclusive environment for neurodivergent creatives.

2.8 Barriers often faced by neurodivergent people include

- * Overstimulating environments (bright lights, loud sounds, crowded areas)
- * Complex forms and application processes.
- * Assumptions around communication styles or social interactions.
- * Inflexible deadlines or schedules

2.9 Tips on increasing access for neurodivergent creatives

- * Use plain jargon free language
- * Offer multiple formats for applications or proposals (Eg. Video, audio, meeting together etc.)
- * Allow flexible deadlines or working times.

*Provide quiet zones at events and in shared work spaces.

* Where possible don't require proof or disclosure of disability to offer access adjustments

*Be conscious of any access docs creative might share with you they can be incredibly helpful for developing long lasting inclusive working relationships.

2.10 Neurodiversity can include:

● **Autism Spectrum Disorder (ASD)**

Autism Spectrum Disorder (ASD) is a lifelong neurodevelopmental condition affecting 1 in 36 children, characterized by persistent challenges in social communication, restricted interests and repetitive behaviors. It is a spectrum with varying support needs often diagnosed by age 2 through behavioral observations while no cure exists, therapies like behavioral, speech and occupational therapy support development and quality of life.

3. Key Aspects of Autism

3.1 Behavioral characteristics & social communication

Social – Emotional Reciprocity: challenges with back and forth conversation, sharing interests or emotions

Non verbal communication : Difficulty with eye contact body language and understanding gestures. *Relationships:* challenges in developing, maintaining and understanding relationships including difficulty in adjusting behavior to social contexts.

Repetitive Behaviors: Engaging in repetitive movements (eg. hand flapping, rocking) lining up toys or using speech in repetitive ways.

Routine & interests: strong need for predictability resistance to change and intense focused interests Sensory processing- heightened or reduced sensitivity to sounds lights or smells.

Types and Spectrum

Previously, different diagnoses existed (eg. Aspergers syndrom) but they are now all classified under the umbrella of Autism Spectrum Disorder(ASD)

Diagnosis:

- Diagnosis is based on behavioral observation and developmental history
- Signs can often be detected by 12-18 months of age with formal consistent diagnosis usually occurring by age 2 or 3.
- It requires a professional comprehensive evaluation

3.2 Causes and prevalence

- Autism is a brain based developmental disability.
- It is believe to be caused by a combination of genetic and environmental factors.
- It is not caused by vaccines
- it is estimated that in 1 in 36 children(or up to 1 in 31 in some studies) are identified with ASD with a higher prevalence in males.

3.3 Treatment and support

There is no cure, but early specialized support can improve abilities. **Therapies:** Behavioral, Speech- language, and occupational therapies. **Support:** Educational and sometimes medication to manage specific symptoms. **ADHD: (Attention – Deficient Hyperactivity)**

One of the most common mental disorder affecting children. It is a chronic neurodevelopmental disorder characterized by persistent in attention, hyperactivity and impulsivity that interfere with daily functioning. It often begins in childhood and lasts into adulthood. Treatment involves a combination of behavioral therapy, medication (stimulants or non-stimulants) and support strategies to manage symptoms effectively.

3.4 Symptoms of ADHD

Symptoms generally fall into three categories

Inattention: Difficult staying organized sustaining focus following instructions and finishing tasks/ Hyperactivity; Excessive restlessness, talking excessively and inability to stay seated.

Impulsivity: Interesting others , blurting out answers and having trouble waiting for a turn.

3.5 Causes and risk factors.

While the exact cause is unknown, ADHD is considered a brain based condition with strong links to genetics and brain chemistry or structure.

Environmental factors like premature birth, low birth weight, or exposure to lead or toxins during pregnancy or early childhood.

3.6 Treatment Approaches

Behavioral Therapy: Training for parents, classroom management and social skills training are primary for children.

Medication: Stimulant medications (eg. Adderall, Ritalin) are common along with non stimulant alternatives

support: A combination of education therapy and structure.

ADHD is not caused by laziness but rather an intense often exhausting struggle to overcome cognitive barriers

Symptoms of ADHD

Common symptoms of ADHD include inattention (difficulty focusing, making careless mistakes, disorganization), hyperactivity (fidgeting, restlessness, excessive talking), impulsivity (blurting answers, interrupting, trouble waiting turns), forgetfulness and difficulty in completing tasks.

All of which impact daily functioning and self-regulation. ADHD is characterized by Inattention, Hyperactivity and Impulsivity. It is not laziness but a neurological difference affecting executive functioning. Usual treatment for ADHD are behavioral therapy, medication, and structured support.

Learning Difficulties

- Dyslexia (Reading)
- Dyspraxia (coordination)
- Dyscalculia (math)
- Tourette Syndrome

Neurodivergent individuals frequently experience:

- Chronic anxiety and depression
- Burnout from masking (suppressing traits to appear neurotypical)
- Sensory overload
- Social isolation
- Low self-esteem
- Misdiagnosis or diagnostic overshadowing

4. Disability Studies and Mental Health

Disability studies challenge the traditional medical model of disability. It focuses instead on social, cultural and political factors that shape the experience of disability and mental health conditions. This field explores the high prevalence of mental health struggles among people with the disabilities given by factors like Stigma, reduced social integration and environmental barriers.

4.1 Core Concepts and Intersection

Mad studies: An interdisciplinary field often intersecting with disabilities studies that challenges traditional psychiatric narratives and centers the expertise of those labeled with mental illness or “Madness”.

4.2 Social model of disability: Emphasis that disability is not just an individual impairment but a result of social, environmental and attitudinal barriers including ableism.

Intersectional Approach: Analyzes how mental health challenges are often compounded by other forms of oppression or social marginalization.

4.3 Key Trends and Issues

High prevalence: People with disabilities have significantly higher odds of experiencing anxiety and depression compared to the general population. **Stigma and Barriers:** Stigma, discrimination and inadequate access to care worsen mental health outcomes for individuals with disabilities.

4.4 Interaction with caregivers:

Parents and caregivers of children with disabilities also face elevated risks of poor mental health and increased service utilization.

Cultural context: Mental health and disability are often culturally constructed, requiring critical analysis of how these concepts are understood and treated in society.

5. Research and Actionable Areas

Policy Improvement: Integrating Mental Health Promotion into broader disabilities support services and vice versa.

Inclusive services: Addressing ableism in mental health care settings to improve the quality of care for disabled individuals.

Self Advocacy: Prioritizing the voices and experience of people with lived experiences of both disability and mental health issues.

5.1 Mental Health Implications of living with Disability

Living with disability significantly increases the risk of mental health issues, with studies showing disabled adults are up to 5 times more likely to experience frequent mental distress than their non-disabled peers. Common implications include high rates of depression, anxiety, social isolation and chronic stress, often driven by societal stigma, physical pain and reduced independence.

6. Key Mental Health implications

Depression and Anxiety: Individuals with disabilities face a significantly higher prevalence of depression and anxiety with the some studies showing roughly 70% exhibit depression symptoms and 55% report anxiety.

Frequent Mental Distress: Over 30% of adults with disabilities experience frequent mental distress. *Social isolation and loneliness:* Barriers to accessibility, work and education often lead to reduced social integration, causing feelings of isolation.

Chronic pain and fatigue: Physical limitations and pain decreased energy and mental fatigue *Identity and self-esteem issues:* Low self esteem and identity disturbance(diffusion) . *Impact of stigma:* Societal bias and discrimination can cause significant emotional distress .

6.1 Mental Health Accessibility and inclusion barriers:

Mental health accessibility for people with disabilities is hinder by systemic attitudinal and physical barriers leading to higher rates of untreated distress. Key obstacles include a shortage of disability trained professionals inaccessible facilities and digital tools. Financial burdens and deep seated social stigma. These issues often cause social isolation worsening conditions like anxiety and depression.

6.2 Key accessibility and inclusion barriers

Attitudinal and societal stigma : Prevalent biases including infantilization and lack of understanding from providers lead to the invalidation of mental health concerns nearly 70% of people with disabilities (PWD) report anticipated provider bias as a barrier to care.

Lack of trained professionals: Many mental health providers lack training to work with PWD, causing them to feel ill - equipped or to mistakenly attribute mental health issues solely to the disability.

Systemic & Economic obstacles: Financial strain from medical costs and lower employment rates create barriers to accessing therapy.

Physical and Digital Inaccessibility: Inaccessible buildings, lack of transportation and in some cases inaccessible telehealth platforms prevent individuals from receiving care.

Societal isolation and exclusion: barriers to participate in community life, education and work create profound loneliness and consequently severe depression or anxiety.

6.3 Impact on Mental health: PWD are at a greater risk of poor mental health due to the intersectional challenges. The World Health Organization (WHO) notes that ableism and discrimination in daily life and health care settings aggravate mental health conditions with social exclusion being a primary driver of poor well-being.

Social and medical model of disability : Social model of disability says that disability is caused by the way society is organized ,rather than by a person's impairment or differences it looks at ways of removing barriers that restrict life choices for disabled people. When barriers are removed disabled people can be independent and equal in society, with choice and control over their own lives.

Disabled people developed the social model of disability because of the traditional medical model did not explain their person experience of disability or help to develop more inclusive ways of living. *Medical model of disability:* It says that people are disabled by their impairments or differences under the medical model, these impairments or differences should be fixed or change by medical and other treatments even when the impairment or difference does not cause pain or illness.

The medical model looks at what is 'wrong' with the person and not what the person needs. It creates low expectations and leads to people losing independence, choice and control in their own lives.

6.4 Social model of disability some examples:

* *A wheelchair user wants to get into a building with a step at the entrance.* Under a social model solution, a ramp would be added to the entrance so that the wheelchair user is free to go into the building immediately .Using the medical model, there are very few solutions to help wheelchair users to climb stairs, which excludes them from many essential and leisure activities.

* *The teenager with a learning disability wants to work towards living independently in their own home but is unsure how to pay the rent,* under the social model the person would be supported so that they are unable to pay rent and live in their own home. Under a medical model, the young person might be expected to live in a communal home.

* *A child with a visual impairment wants to read the latest best-selling book to chat about with their sighted friends.* Under the medical model there are very few solutions but a social model solution ensures full text audio recordings are available when the book is first published. This means children with visual l impairments can join in with cultural activities on and equal basis with everyone else. **Intersectional challenges:** Intersectional challenges in mental health arises when multiple social identities such as race, gender, class, sexuality and ability overlap, creating unique compounder systems of discrimination and inequality. These intersections often involving racism, sexism or ableism lead to higher rates of anxiety, depression and PTSD while creating barriers to culturally competent care.

6.5 Compounded stigma and discrimination : Individuals holding multiple marginalized identities (eg. the queer person of colour or a disabled person from a lower socio economic background face layers of prejudice that exacerbate mental health issues.

6.6 Barriers to care: Marginalized groups often experience reduced access to mental health services due to financial constraints, geographical or language barriers.

Cultural misalignments: Mental health services often lack cultural competence failing to understand the unique experience of diverse populations which results in improper diagnosis or treatment.

6.7 Chronic stress and trauma: The constant experience of micro aggressions, systemic inequalities in prejudice acts as a chronic stresser, significantly impacting mental well being.

6.8 Unique group challenges: Specific intersections create distinct crisis such as the mental health consequences of navigating rigid gender norms alongside racial discrimination.

Addressing these challenges requires a shift from viewing mental health as a uniform experience to acknowledging the specific layered and contextual realities of each individual.

Culturally competent care, policy changes and awareness of systemic oppression are essential for improving mental health outcomes.

6.9 Effects of Gender neurodivergence and disability in mental health

The intersections of gender neurodivergence and disability creates unique compounded and often severe impacts on mental health driven by structural, societal and systemic inequalities. Individuals at the intersections of these identities often face due to the pressure of 'mask' or uniform to neurotypical or asnormative standards in a society not designed for them.

6.10 Effects of Neurodivergence on Mental Health

High Co- occurrence: Neurodivergent individuals (autistic, ADHD, etc) are significantly more likely to experience mental health issues compared to neurotypical peers, with up to half of autistic adults experiencing depression and half of ADHD adults facing anxiety disorders.

Masking and Burnout: Forcing oneself to hide neurodivergent traits (masking/ camouflaging) to fit in causes severe long term mental strain, leading to burnout, low self worth and extreme anxiety.

Sensory and communication overload: Navigating a world designed for neurotypical brains can result in chronic stress meltdowns and social isolation.

Delayed Diagnosis: Particularly for women and girls, a lack of timely diagnosis leads to years of feeling unsupported, misunderstood and misdiagnosed, resulting in higher risks of mental health conditions.

Effects of Gender on Mental Health- Intersecting with Neurodivergence

Higher Risks for women and Non- binary individuals: Autistic women and gender-diverse individuals report higher rates of mental ill health (up to 77% diagnosed with a psychiatric condition by age 25) than their male counterparts.

Unique pressures on transgender individuals: Transcenders and gender-diverse (TGD) people face higher rates of anxiety, depression substance use and suicide due to stigma and discrimination.

Misdiagnosis of women: Due to gender bias in diagnostic tools, autistic women are frequently miss diagnosed with conditions like borderline personality disorder or eating disorders instead of receiving an autism diagnosis. Female Orphan present with internalizing behaviours anxiety.

Gendered symptom expression: Female often present with internalizing behaviours (anxiety, depression) which are often missed or misinterpreted by clinicians compared to the externalizing behaviours (disruptive, impulsive) more common and males.

7. Conclusion

Mental health disparities cannot be understood without examining the intersections of gender, neurodivergence, and disability. These identities do not operate independently, they overlap to create compounded vulnerability within systems shaped by stigma, structural inequality, and exclusion. To move forward, we must reduce stigma through public education, expand inclusive and affirmative care, integrate disability and mental health services, reform diagnostic tools to address gender bias and center lived experiences in policy-making etc. Mental health justice requires more than treatment — it requires transformation of the systems that produce inequality.

References

1. Harari, Yuval Noah. *Sapiens: A Brief History of Humankind*. Harvill Secker, 2014.
2. Phadke, Shilpa, et al., editors. *Disability and the Postcolonial*. Routledge, 2020.
3. Singer, Judy, and Alison Broadhurst, editors. *The Neurodivergent Reader*. PCCS Books, 2020.
4. Taylor, Sonya Renee. *The Body is Not an Apology: The Power of Radical Self-Love*. Berrett-Koehler Publishers, 2021.
5. Butler, Judith. *Gender Trouble: Feminism and the Subversion of Identity*. Routledge, 1990.
6. Murphy, Joseph. *The Power of Your Subconscious Mind*. Prentice Hall, 1963.
7. *Autism in Adulthood*. Mary Ann Liebert, Inc., ISSN 2573-9582.
8. *Disability & Society*. Taylor & Francis Online, ISSN 0968-7599.
9. *International Journal of Disability and Social Justice*. Open-access, ISSN 2634-5255.
10. *Journal of Gender Studies*. Taylor & Francis Online, ISSN 0958-9236.
11. *Journal of Neurodiversity*. Open-access journal, ISSN 2634-5263. Articles
12. "The intersectionality of gender and disability." *Disability & Society*, vol. 33, no. 5, 2018, pp. 663-678. Taylor & Francis Online.
13. "Neurodiversity and mental health." *Journal of Neurodiversity*, vol. 1, no. 1, 2020, pp. 1-10. Open-access.
14. "Gender and mental health disparities." *Journal of Gender Studies*, vol. 28, no. 3, 2019, pp. 257-272. Taylor & Francis Online.